

DOCUMENT RESUME

ED 326 734

CG 022 480

AUTHOR Newman, Isadore; And Others
TITLE Drugs, Suicide, and Rock and Roll.
PUB DATE Apr 90
NOTE 40p.; Paper presented at the Annual Meeting of the Ohio Academy of Science (99th, Dayton, OH, April 1990).
PUB TYPE Speeches/Conference Papers (150)
EDRS PRICE MF01/PC02 Plus Postage.
DESCRIPTORS *Adolescents; *Drug Use; *Locus of Control; *Music; Parent Child Relationship; Peer Relationship; Religion; *Suicide
IDENTIFIERS *Lyrics

ABSTRACT

This study was conducted to investigate the relationship of music preference to locus of control, religiosity, parent and peer relationships, and drug use, with an emphasis on the relationship between musical preference and suicide ideology. Adolescent clients (N=22) at a large Christian counseling center and adolescents (N=76) at a religious after-school program completed an instrument measuring the constructs of suicidal risk, social support, family and peer relationships, and drug involvement. The results for the non-clinical sample indicated a negative relationship between preference for rock music and drug involvement and suicide, and a positive relationship between parent relationship and preference for rock music. Unknown music preference was negatively related to parent relationship. The findings revealed that music preference was not significantly related to most of the subscales in the non-clinical sample. Overall music preference did not differentiate between the clinical and non-clinical groups. For the clinical group, preference for heavy metal music was related to suicide ideology and drug involvement. These findings may suggest that while non-clinical adolescents are not as affected by the messages conveyed in heavy metal music, clinical adolescents may be more sensitive to the messages and mood of this type of music. (NB)

* Reproductions supplied by EDRS are the best that can be made *
* from the original document. *

DRUGS, SUICIDE, AND ROCK AND ROLL

I. Newman, G. J. Marchant, M. L. Newman,
G. Gertz, and R. McMorrow

April, 1990

Running head: Suicide and Rock

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

- ☒ This document has been reproduced as
received from the person or organization
originating it.
☐ Minor changes have been made to improve
reproduction quality.

- Points of view or opinions stated in this docu-
ment do not necessarily represent official
DERI position or policy.

PERMISSION TO REPRODUCE THIS
MATERIAL HAS BEEN GRANTED BY

Isadore Newman

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC).

A paper presented at the 99th annual meeting of the Ohio Academy
of Science (Psychology Division), Dayton, Ohio.

DRUGS, SUICIDE, AND ROCK AND ROLL

I. Newman, G. J. Marchant, M. L. Newman, G. Gertz, & R. McMorrow

Adolescents are at risk in regard to suicide and that risk is increasing (Berman, 1984). The number of reported suicide deaths are believed to be highly conservative estimates of the problem. Some estimates indicate that there are six suicide attempts for every one successful completion. Other estimates are that there are between 50 - 150 attempts for every completion (Wodarski, 1989). Attempted suicides seem to be cries for help by adolescents who perceive their problems as insurmountable. Suicide is the third leading cause of death in the 15 to 20 year old age group (U.S. Bureau of Vital Statistics, 1978). This is a 1978 statistic and we know that teenage suicide has increased 237% between 1960 and 1980.

Parents of adolescents, educators, and mental health professionals have been concerned about the potential effects of the explicit lyrics in popular music, as possibly facilitating or effecting the attitudes and/or behaviors that lead to suicide and drug use. The National Parent Teachers Association (PTA) and the Parent's Music Resource Center have disseminated to its members, their concerns about explicit lyrics and themes in some of the most popular teen music. A variety of studies (eg. Prinsky & Rosenbaum, 1987) have indicated that teenagers do not understand the lyrics or the message in most of their favorite rock music.

However, there are a variety of methodological problems in their study as well as a conceptual one. Teenagers may not get the message at a cognitive level and yet may receive it at an emotional level.

More recent research by Yee, Britten, and Thomsom (1988), found that adolescents who prefer heavy metal music were significantly more likely to favor sadism, drug and alcohol use, and were more likely to participate in premarital sex as well as have a significantly more negative attitude toward school. Interestingly, there was also a positive correlation between listening to heavy metal music and viewing x-rated movies and reading pornographic magazines. However, these researchers concluded that the variance accounted for was small, ranging between a high of 10% and a low of 3%.

The purpose of this study was to investigate the relationship of music preference as it relates to locus of control, religiosity, parent and peer relationships, and drug use, with an emphasis on the relationship between musical preference and suicide ideology. It is believed that locus of control is an important mediating variable in understanding the potential relationships between music preference, drug involvement, peer relationships, family relations, and suicide ideology.

Method

Subjects

Twenty-two subjects were from a large Christian counseling center which asked their adolescent clients between the ages of 13-18, to voluntarily complete an instrument which measured the constructs of suicidal risk, social support, family and peer relationships and drug involvement. These subjects were also rated by their counselors on a five-point scale as being ranging from low-risk to high-risk for suicide. An additional 76 subjects, ages 14-17, at a religious after-school program, were asked to voluntarily fill out the same questionnaire. These however, did not have a therapist's clinical ratings of their suicide risk. The total sample included 98 subjects.

Instrument

The instrument used is a 46 item questionnaire, measuring the following seven concepts (See Appendix A for a copy of the questionnaire.):

1. Suicide ideology - (19 items) - Nineteen items, some of which had reverse scoring, measured suicide ideology. High scores on items 1, 2, 13, 20, 25, 33, 37, and 41 were perceived as low suicide ideology. High scores on items 3, 19, 21, 27, 30, 35, 39, 40, 43, 44, and 45 were perceived as a high suicide ideology score.

2. Locus of control - (4 items) - Items 4-7 were used to estimate locus of control. The higher the score, the more external the locus of control. These items have previously been identified by Rotter as the best four items of his scale, and were incorporated in the Coleman's High School and Beyond data.
3. Peer relationships - (5 items) - Five items measured peer relationships. A high score on items 8, 22, and 29 were positively scored and a high score on items 12 and 16 were negatively scored.
4. Social support - (1 item) - Only one item, number 9, was used to measure social support. Previous research indicated that the larger the support network (the number of people who could be identified, regardless of quality), the larger the perceived social support. Therefore, this one item was used to measure social support.
5. Religiosity - (5 items) - Five items were used to measure religiosity. One item, number 26, was not used in measuring the total score, since it could not be determined if a high score was positive or negative. For items 10, 14, 17, and 23 high scores were positive.
6. Parent relationships - (8 items) - Eight items were used to estimate parent relationships. On items 11, 24, 28, 31, and 38, high scores were positive ratings. For items 15, 18 and 34, high scores were negative

ratings.

7. Drug involvement - (4 items) - Four items, 32, 36, 42, and 46, measured drug involvement. High scores indicated high involvement in drug use.

Results

The frequencies and demographic information were presented for the clinical and nonclinical groups (see Tables 1 and 2). There appears to be more similarities than differences between the two groups. The means and standard deviations for the subscales and factors were calculated for the groups (see Tables 3 and 4). The correlation matrix for the nonclinical group for music preference and the subscales and factors indicated a few significant relationships (see Table 5). There was a negative relationship between preference for rock and drug involvement and suicide, and a positive relationship between parent relationship and preference for rock. Unknown music preference was negatively related to parent relationship. For the clinical sample there were positive correlations between preference for heavy metal and suicide and drug involvement (see Table 6). The music preference categories did not differentiate between the clinical and nonclinical groups (see Table 7).

Typologies were estimated through a Q-factor analytic procedure run separately on the clinical and nonclinical samples. The Q-factor analysis on the clinical sample indicated a two

factor solution (Table 8a-d). Of the 22 people in the clinical group, 13 were identified as type 1 and 6 were identified as type 2. Type 2 people tended to score higher on items which indicated more positive relationships with family and friends. The Q-factor analysis on the non-clinical group also produced a two factor solution. With 47 people in type 1 and 21 people in type 2. Both of the non-clinical typologies would be considered positive in nature. The type 2 group tended to have better parent relations, were more concerned with personal appearance, and were the lowest on suicide ideology.

Discussion

One of the most interesting features of the results was that music preference was not significantly related to most of the subscales in the non-clinical sample. Overall music preference did not differentiate between the clinical and non-clinical groups. However for the clinical sample, preference for heavy metal music was related to suicide ideology and drug involvement. One possible implication is that "non-clinical" adolescents are not as affected by the messages conveyed in heavy metal music. However, "clinical" adolescents may be more sensitive to the messages and mood of this type of music. Since this was a correlational study an equally valid argument can be made which indicates that "clinical" adolescents are more likely to be attracted to the mood and message of heavy metal music. However,

the relative similarity of music preferences between clinical and non-clinical groups did not suggest this. This is somewhat consistent with recent literature which indicates that there is a very small effect or no effect of music preference to pathology in adolescents, contrary to popular expectations.

This study also investigated typologies of "clinical" and "non-clinical" adolescents. Further research needs to be directed toward the relationship of the typologies to music preference and pathology. It is this approach to typologies which may provide insights into the characteristics of adolescents. These characteristics may provide direction for diagnosis as well as prevention strategies.

It is important to recognize the nature and limitations of this study, however, it is equally important that the relationships described here receive further investigation considering the nature of the topic.

Table 1

Frequencies and Means of Demographics
for Nonclinical Group

Males	40
Female	36
Average Age	13.86
Average Grade Level	8.1
Average Number Siblings	1.4
Like Heavy Metal Music	33%
Like Western Music	2%
Like Jazz Music	21%
Like Pop Music	47%
Like Rock Music	80%
Like Classical Music	17%
Like New Wave Music	38%
Unknown	3%
Don't Listen to Music	2%

Table 2

Frequencies and Means of Demographics
for Clinical Populations

Males	5
Female	17
Average Age	15.4
Average Grade Level	9.5
Average Number Siblings	1.8
Like Heavy Metal Music	45%
Like Western Music	0%
Like Jazz Music	4%
Like Pop Music	45%
Like Rock Music	68%
Like Classical Music	27%
Like New Wave Music	18%
Christian Rock	27%
Don't Listen to Music	4%

Table 3

Means and SD for the Clinical Group

	Means	SD
Suicide	39.25	11.55
Locus	8.13	2.39
Friends	15.95	4.17
Religion	14.88	2.19
Parents	25.21	4.17
Drugs	6.31	3.06
Support	16.54	23.26
Factor I	45.27	5.39
Factor II	26.72	4.22

Note. $N = 22$

Table 4

Means and SD for the Nonclinical Group

	Means	SD
Suicide	32.12	8.42
Locus	7.77	1.97
Friends	16.82	2.81
Religion	13.33	3.59
Parents	27.92	3.51
Drugs	5.92	2.74
Support	14.43	17.42
Factor I	45.79	3.90
Factor II	22.98	3.50

Note. N = 76

Table 5

Correlation of Music Preferences with Scales
and Factor Scores for the Nonclinical Population

		Heavy Metal	Western	Jazz	Pop	Rock	Classical	New Wave	Unknown	Do Not List
Suicide	r =					-.25				
	p =					.03				
Locus	r =									-.2
	p =									.0
Friends	r =									
	p =									
Religion	r =									
	p =									
Parents	r =					.36			.33	
	p =					.002			.007	
Drugs	r =					-.38				
	p =					.0009				
Support	r =	.29			.22					
	p =	.009			.05					
Factor I	r =		-.29							
	p =		.01							
Factor II	r =									
	p =									

Note. Correlations with p values of .05 or less are reported.

Table 6

Correlation of Music Preferences with Scales
and Factor Scores for the Clinical Population

		Heavy Metal	Western	Jazz	Pop	Rock	Classical	New Wave	Christian Rock	Don't Listen
Suicide	r =	.5								
	p =	.02								
Locus	r =									
	p =									
Friends	r =									
	p =									
Religion	r =									
	p =									
Parents	r =									
	p =									
Drugs	r =	.54								
	p =	.008								
Support	r =									
	p =						.4			
							.06			
Factor I	r =									
	p =									
Factor II	r =									.57
	p =									.005

Note. Correlations with p values of .05 or less are reported.

Table 7

Correlation Between the Nine Categories of Music
with the Clinical vs. Nonclinical group

		Clinical/Nonclinical
Heavy	r =	NO
Metal	p =	
Western	r =	SIGNIFICANT
	p =	
Jazz	r =	CORRELATIONS
	p =	
Pop	r =	
	p =	
Rock	r =	
	p =	
Classical	r =	
	p =	
New Wave	r =	
	p =	
Christian	r =	
Rock	p =	
Don't	r =	
Listen	p =	

Note. Only scores with p values of .05 or less will be entered.

Table 8a

VARIABLE ASSIGNMENTS WITH FACTOR WEIGHTS BY TYPE

VAR.	WEIGHT
TYPE 1 (N= 13)	
2	2.7853
3	5.6623
7	3.6819
9	7.1501
10	1.9744
12	0.6428
14	3.2261
15	6.6756
16	8.5488
18	5.6635
19	5.8715
21	13.1856
22	1.8630
TYPE 2 (N= 6)	
1	0.7142
4	3.7355
6	3.2116
8	0.8844
13	1.0996
17	0.5146
TYPE 3 (N= 3)	
5	2.1473
11	0.4876
20	1.1318

Note. Table 8 is for the clinical sample.

DESCENDING ARRAY OF Z-SCORES AND ITEM DESCRIPTIONS FOR TYPE 1

16

DESCENDING ARRAY OF Z-SCORES AND ITEM DESCRIPTIONS FOR TYPE 2

[illegible]

Table 8d

ITEM DESCRIPTIONS AND DESCENDING ARRAY OF DIFFERENCES BETWEEN TYPES 1 AND 2

	1	2	DIFFERENCE	
9.	6.565	-1.719	8.284	9. How many people can you name
16.	-0.411	-1.390	0.979	that would be willing to
15.	-0.441	-1.347	0.9	help you if you were in need?
21.	-0.341	-1.208	0.867	
26.	-0.198	-1.053	0.855	
30.	-0.323	-1.157	0.834	
36.	-0.380	-1.194	0.814	16. People my age don't like me.
19.	-0.441	-1.245	0.804	
35.	-0.325	-1.129	0.804	15. My parents/guardians don't
42.	-0.447	-1.245	0.798	expect anything of me .
46.	-0.467	-1.245	0.778	
34.	-0.208	-0.927	0.719	
40.	-0.341	-0.992	0.651	21. I think of things too
4.	-0.426	-0.977	0.551	bad to tell other people .
27.	-0.264	-0.725	0.461	
32.	-0.225	-0.488	0.263	
39.	-0.192	-0.442	0.251	
12.	-0.332	-0.509	0.177	
13.	-0.025	-0.200	0.174	
43.	-0.388	-0.535	0.147	
45.	-0.143	-0.233	0.090	
44.	-0.129	-0.215	0.087	
6.	-0.284	-0.290	0.006	
5.	-0.186	-0.073	-0.113	
2.	0.139	0.333	-0.194	
23.	0.090	0.418	-0.328	
33.	0.049	0.472	-0.423	
18.	-0.064	0.365	-0.429	
31.	-0.260	0.266	-0.526	
41.	0.034	0.742	-0.708	
10.	-0.186	0.532	-0.717	
7.	-0.205	0.521	-0.726	
22.	0.156	0.974	-0.818	
3.	-0.233	0.635	-0.867	
20.	0.054	0.931	-0.878	
1.	0.130	1.053	-0.923	
14.	-0.196	0.892	-1.088	
17.	0.004	1.105	-1.102	
38.	-0.173	0.935	-1.108	
8.	0.114	1.343	-1.229	
29.	0.113	1.352	-1.239	
11.	0.078	1.319	-1.241	
37.	0.227	1.495	-1.268	
25.	0.206	1.619	-1.413	
28.	0.143	1.619	-1.476	37. I care about the way I look.
24.	0.133	1.619	-1.486	28. I love my parents/guardians.
				24. My parents/guardians love me.
				25. I feel many people care for me.

FACTOR 3 IS 43.54 PER CENT NEGATIVE.

TABLE 9a

NEGATIVE ITEMS ARE EXTRACTED, MADE POSITIVE AND FORMED INTO ADDED TYPE 6
FACTOR 4 IS 40.14 PER CENT NEGATIVE.

NEGATIVE ITEMS ARE EXTRACTED, MADE POSITIVE AND FORMED INTO ADDED TYPE 7
FACTOR 5 IS 36.10 PER CENT NEGATIVE.

NEGATIVE ITEMS ARE EXTRACTED, MADE POSITIVE AND FORMED INTO ADDED TYPE 8

VARIABLE ASSIGNMENTS WITH FACTOR WEIGHTS BY TYPE

VAR.	HEIGHT
TYPE 1 (N= 47)	
2	3.4975
5	41.1974
6	2.8694
7	19.4614
8	0.8033
9	1.8520
11	1.2441
12	3.9978
13	16.1990
14	20.3443
15	2.5244
17	11.3913
18	10.6856
19	1.8184
20	1.8782
22	10.0823
23	10.7869
26	3.3058
27	10.4041
28	1.7331
29	32.5601
30	19.8266
37	3.2442
38	4.8009
39	15.3015
40	8.3173
41	3.5060
43	11.7008
44	13.9516
45	14.3328
46	2.8601
47	2.9524
49	8.7439
50	4.4439
52	10.8902
53	2.9525
54	15.6759
55	9.8799
56	5.5161
59	3.6962
60	5.0276
61	1.5735
62	2.3738
63	0.5320
65	2.3062
67	4.0160
74	4.9704

Table 9b

TYPE 2 (N= 21)	
1	2.5245
3	1.4385
4	2.5690
16	2.7246
21	3.1704
25	1.8922
31	0.8647
32	1.4240
35	0.5812
36	0.4404
42	0.5347
48	1.8376
51	1.9444
57	2.4701
58	0.7633
68	0.9370
69	3.1684
70	3.2154
71	2.2109
72	1.9879
73	1.9542
TYPE 3 (N= 4)	
24	0.6292
33	0.4291
34	1.0202
66	0.7240
TYPE 4 (N= 1)	
64	0.2345
TYPE 5 (N= 1)	
10	0.3288
TYPE 6 (N= 0)	
TYPE 7 (N= 0)	
TYPE 8 (N= 0)	

NO SUBJECTS LOAD HIGHEST ON TYPE(S) 6 7 8
THOSE TYPES ARE ELIMINATED.

Note. Table 9 is for the clinical sample.

Table 9c

DESCENDING ARRAY OF Z-SCORES AND ITEM DESCRIPTIONS FOR TYPE 2

[illegible]

37. I care about the way I look.

28. I love my parents/guardians.

11. My parents/guardians value me as a person

19. I think of killing myself

46. I sometimes get drunk or high alone.

9. How many people can you name that would be willing to help you if you were in need?

Table 9d

DESCENDING ARRAY OF Z-SCORES AND ITEM DESCRIPTIONS FOR TYPE 1

[illegible]

9. How many people can you name that would help you if you were in need?

28. I love my parents/guardians.

24. My parents/guardians love me. .

25. I feel many people care for me.

46. I sometimes get drunk or high alone.

42. I often use alcohol and/or drugs.

43. I have thoughts of how to do myself in.

Table 9e

ITEM DESCRIPTIONS AND DESCENDING ARRAY OF DIFFERENCES BETWEEN TYPES 1 AND 2

	1	2	DIFFERENCE
9.	6.458	-2.176	8.634
46.	-0.491	-1.253	0.762
19.	-0.474	-1.211	0.737
30.	-0.424	-1.156	0.732
43.	-0.496	-1.187	0.690
42.	-0.495	-1.081	0.586
40.	-0.373	-0.943	0.570
16.	-0.411	-0.970	0.559
15.	-0.424	-0.903	0.479
36.	-0.471	-0.895	0.425
35.	-0.475	-0.864	0.388
7.	-0.117	-0.503	0.386
26.	-0.488	-0.861	0.372
34.	-0.465	-0.831	0.366
4.	-0.377	-0.718	0.340
39.	-0.369	-0.707	0.338
32.	-0.352	-0.666	0.314
44.	-0.288	-0.595	0.307
23.	-0.222	-0.479	0.257
27.	-0.208	-0.430	0.222
6.	-0.301	-0.490	0.188
3.	-0.226	-0.407	0.181
18.	-0.220	-0.357	0.137
21.	-0.323	-0.367	0.044
45.	-0.260	-0.271	0.011
5.	-0.349	-0.304	-0.044
12.	-0.308	-0.072	-0.236
17.	0.001	0.254	-0.254
14.	-0.039	0.371	-0.410
10.	0.062	0.609	-0.548
22.	0.174	0.758	-0.584
8.	0.229	0.930	-0.701
29.	0.135	0.852	-0.717
2.	0.158	0.967	-0.809
31.	0.076	0.909	-0.833
33.	0.158	1.141	-0.982
13.	0.103	1.164	-1.061
1.	0.238	1.311	-1.073
41.	0.105	1.205	-1.099
20.	0.214	1.318	-1.104
38.	0.159	1.278	-1.120
25.	0.240	1.428	-1.188
24.	0.261	1.452	-1.191
11.	0.213	1.519	-1.306
28.	0.247	1.613	-1.366
37.	0.219	1.620	-1.400

Bibliography

- American Humane Association. (1989). Highlights of official aggregate child neglect and abuse reporting, 1986. Author. A sharp tongue can inflict lasting damage. (1989, March 7). USA Today.
- Anderson, D. B., & McClean, L. J. (1960). Identifying suicide potential. New York: Behavioral Publications.
- Asarnow, J. R., Carlson, G. A., & Guthrie, D. (1987). Coping strategies, perception, hopelessness, and perceived family environments in depressed and suicidal children. Journal of Consulting and Clinical Psychology, 55 (3), 361-366.
- Barrell, T. (1984). A friend in need. Ladies Home Journal, 7, 18-21.
- Barrett, T. C. (1985). Youth in crisis: Seeking solutions to self-destructive behavior. Longmont, CO: Sopris West.
- Barrett, T. C. (1989). Teens in crisis. Arlington, VA: American Association of School Administrators.
- Barrett, T. C., & Barrett, M. (1989). When tragedy strikes. Longmont, CO: Sopris West.
- Bassuk, E. L. (1982). Lifelines, clinical perspectives on suicide. New York: Publishing Corporation.
- Bellak, L., & Seigel, H. (1983). Handbook of brief and emergency psychotherapy. Larchmont, NY: CPS.
- Best, J., & Luckenbill, D. F. (1982). Organizing behavior. Englewood Cliffs, NJ: Prentice-Hall, Inc.
- Bhatia, S. C., Khan, M. H., & Sharma, A. (1986). Suicide risk: Evaluation and management. American Family Physician, 34 (3), 167-174.
- Bleiberg, E. (1988). Adolescence, sense of self, and narcissistic vulnerability. Bulletin of the Menninger Clinic, 52 (3), 211-229.
- Brain, L. A. (1985). Deadly cry for help. Engage, 2, 94, 13-17.

- Brophy, J. (1987, October). Developing a youth suicide prevention program. Thrust, pp. 40-41.
- Byron, P. (1989, January 8). Recognizing depression in teens. The Fresno Bee, p. 1, (B).
- Chance, S. (1988). Surviving suicide: A journey to resolution. Bulletin of the Menninger Clinic, 52(1), 30-39.
- Clark, A., Clemes, H., & Bean, R. (1980). How to raise teenagers' self-esteem. Enrich the Whole Child series. Sunnyvale, CA: Enrich and Ohaus.
- Coleman, L. (1987). Today's epidemic of youth suicide clusters. In Suicide clusters. New York: Faber & Faber.
- Coleman, L. (1987). Suicide warning signs. In Suicide clusters. New York: Faber & Faber, Inc.
- Coleman, L. (1987). Teen suicide checklist. In Suicide clusters. New York: Faber & Faber, Inc.
- Cousart, F. (1988, September 23). Fresno Unified School District will try to revive peer counsel program. The Fresno Bee, 1, (B).
- Crow, G. A., & Crow, L. I. (1987). Crisis intervention and suicide prevention. Springfield, IL: Charles C. Thomas.
- Douglas, L. (1986). Is adolescent suicide a third degree game and who is the real victim? Transactional Analysis Journal, 16 (3), 165-169.
- Dunne, E. J., McIntosh, J. L., & Dunne-Maxim, K. (1987). Suicide and its aftermath. New York: W. W. Norton & Co., Ltd.
- Eisenberg, L. (1986). Does bad news about suicide beget bad news? The New England Journal of Medicine, 315 (11), 705-707.
- Fuchs, R. (1985). Presentation of depression in an emergency setting. Behavioral, Cognitive, and Affective Disturbances, 5, 219-232.
- Goleman, D. (1989, January 10). Studies: 1 in 5 children has serious emotional troubles. The Fresno Bee, p. 8, (A).

- Gould, M. S., & Shaffer, D. (1986). The impact of suicide in television movies. The New England Journal of Medicine, 315 (11), 690-694.
- Haffen, B. Q., & Frandsen, K. J. (1986). Youth suicide. Evergreen, CO: Cordillera Press, Inc.
- Hawton, K., (1986). Suicide and attempted suicide among children and adolescents. Beverly Hills: Sage.
- Hawton, K., & Catalan, J. (1982). Attempted suicide: A practical guide to its nature and management. New York: Oxford University Press.
- Hendin, H. (1982). Suicide among the young. In Suicide in America. New York: W. W. Norton, Inc.
- Hendin, H. (1985). Suicide among the young: Psychodynamics and demography. In Peck, Farberow and Litman (Eds.), Youth suicide. New York: Springer.
- Hipple, J., & Cimboric, P. (1979). The counselor and suicidal crisis. Springfield, IL: Charles C. Thomas.
- Hubler, E. (1984). Teenage suicide. Ladies Home Journal, 7, 17-20.
- Husain, S. A., & Vandiver, T. (1984). Suicide in children and adolescents. Jamaica, NY: Spectrum Publications, Inc.
- Jacobs, D. (1982-83). Evaluation and care of suicidal behavior in emergency settings. International Journal of Psychiatry in Medicine, 12 (4), 295-310.
- Knopk, I. J. (1984). Abnormalities of adolescence and youth. In Childhood psycho-pathology. Englewood Cliffs, NJ: Prentice-Hall, Inc.
- Landau-Stanton, J., & Stanton, D. M. (1985). Treating suicidal adolescents and their families. In Handbook of adolescents and family therapy. New York: Garner Press, Inc.
- Lester, D. (1988). One theory of teen-age suicide. Journal of School Health, 58 (5), 193-194.
- Lester, D. (1987). Suicide as a learned behavior. Springfield, IL: Charles C. Thomas.

- Lester, G., & Lester, D. (1971). Suicide: The gamble with death. Englewood Cliffs, NJ: Prentice-Hall.
- Levine, S. V. (1987). The myths and needs of contemporary youth. Adolescent Psychiatry, 14.
- Miller, D. (1984). Newslink. American Association of Suicidology, 10 (4).
- Mitchell, J. (1979). Adolescent psychology. Toronto: Rinehart and Winston of Canada, Limited.
- Murphy, G. E. (1984). Suicide and attempted suicide. Psychiatry, 1 (71), 1-17.
- Offer, D. (1970). The psychological world of the teenager. New York: Basic Books.
- Peck, M., Farberow, N. L., & Litman, R. E. (1985). Youth Suicide. New York: Springer Publishing Company.
- Perrone, P. A. (1987, September). Counselor response to adolescent suicide. The School Counselor, pp. 51-57.
- Phillips, D. P., & Carstenson, L. L. (1986). Clustering of teenage suicides after television news stories about suicide. The New England Journal of Medicine, 315 (11), 685-689.
- Pretzel, P. W. (1972). Understanding and counseling the suicidal person. Nashville/New York: Abington Press.
- Prinsky, L. E., & Rosenbaum, J. L. (1987). "Leer-ics" or lyrics: Teenage impressions of rock 'n' roll. Youth and Society, 18 (4), 384-397.
- Range, L. M., McDonald, D., & Anderson, H. (1987). Factor structures of Calhoun's youth scale. Journal of Personality Assessment, 51 (2), 262-266.
- Risking the future. (1987). National Academy Press.
- Sanstock, J. W. (1984). The nature of abnormality and problems and disturbances in adolescence. In Adolescence. Dubuque, IA: Wm. C. Brown Publishers.
- Sexually transmitted disease cases reported. (1985). Centers for Disease Control.

- Shneidman, E. S. (1969). Prologue: Fifty-eight years. In E. S. Shneidman (Ed.), On the nature of suicide. San Francisco: Jossey-Bass.
- Shneidman, E. (1987, March). At the point of no return. Psychology Today, pp. 54-58.
- Smith, Jr., C. W., & Bope, E. T. (1986). The suicidal patient. Postgraduate Medicine, 79 (8), 195-202.
- Smith, K., & Crawford, S. (1986, May). Suicidal behavior among "normal" high school students. Suicide and Life Threatening Behavior.
- U.S. Department of Commerce, Bureau of the Census. (no date) School enrollment, social and economic characteristics of students. Current Population Reports, No. 222, 303, 362, 392, 409. Series P-20. Washington, DC: Author.
- U.S. Department of Health and Human Services, Alcohol, Drug Abuse, and Mental Health Administration. (1987). National trends in drug use and related factors among American high school students, 1975-1986. Washington, D.C.: Author.
- Vital statistics of the United States. Washington, D.C.: U.S. Department of Health and Human Services.
- Weiss, M. J. (1984, June). How to stop a suicide. Ladies Home Journal, p. 56.
- West, D. A., & DiVasto, P. V. (1982). Suicide potential. Postgraduate Medicine, 72 (1), 203-212.
- Why school health? (1987). American Association of School Administrators.
- Wordarski, J. S. (no date). Preventive health services for adolescents. Springfield, IL: Charles C. Thomas Publishers.
- Youth Suicide Surveillance. (1986, November).

QUESTIONNAIRE

I.D.# _____ Sex: M F Age: _____ Grade: _____
 Parent/Guardian: _____ Number of Parents: _____
 Number of Siblings: _____ Religious Affiliation: _____
 Check the types of music you like: heavy metal _____ western _____ jazz _____
 pop _____ rock-n-roll _____ classical _____ new age _____ other _____

- | | |
|---|--|
| | Strongly Disagree
Somewhat Disagree
Somewhat Agree
Strongly Agree |
| 1. I look forward to the future | 1 2 3 4 |
| 2. I feel there is a lot of good I can do | 1 2 3 4 |
| 3. I feel tired and run down. | 1 2 3 4 |
| 4. Good luck is more important for success
than hard work. | 1 2 3 4 |
| 5. Every time I try to get ahead, someone or
something stops me. | 1 2 3 4 |
| 6. Planning only makes a person unhappy since
plans hardly ever work out. | 1 2 3 4 |
| 7. People who accept their condition on life are
happier than people who try to change things. | 1 2 3 4 |
| 8. I have a lot of friends | 1 2 3 4 |
| 9. How many people can you name that would be willing to
help you if you were in need? | (No.) |
| 10. Religion is very important to me | 1 2 3 4 5 |
| 11. My parents/guardians value me as a person. | 1 2 3 4 5 |
| 12. I wish I had better friends. | 1 2 3 4 |
| 13. I am doing very well in school | 1 2 3 4 5 |
| 14. I very frequently practice my religion | 1 2 3 4 5 |
| 15. My parents/guardians don't expect anything of me | 1 2 3 4 5 |
| 16. People my age don't like me. | 1 2 3 4 |
| 17. My religious beliefs make me feel good about myself. | 1 2 3 4 5 |
| 18. My parents/guardians expect too much of me | 1 2 3 4 5 |
| 19. I think of killing myself to get back at others. | 1 2 3 4 |

(turn over)

20. I am fun to be with. 1 2 3 4
21. I think of things too bad to tell other people 1 2 3 4
22. I fit in with most people my age. 1 2 3 4
23. My religious beliefs help me know what to do when
I have problems. 1 2 3 4
24. My parents/guardians love me. 1 2 3 4
25. I feel many people care for me. 1 2 3 4
26. I have changed my religious beliefs. 1 2 3 4
27. I have trouble sleeping. 1 2 3 4
28. I love my parents/guardians. 1 2 3 4
29. I am satisfied with my relationship with members
of the opposite sex. 1 2 3 4
30. It is more painful to live than die. 1 2 3 4
31. My parents/guardians trust my decisions. 1 2 3 4
32. Many of my friends get drunk or high. 1 2 3 4
33. I feel very healthy. 1 2 3 4
34. I wish I had different parents/guardians. 1 2 3 4
35. I have thoughts of giving all my things away. 1 2 3 4
36. I sometimes get drunk or high with my friends. 1 2 3 4
37. I care about the way I look. 1 2 3 4 5
38. My mom and dad have a good relationship with
each other. 1 2 3 4 5
39. Most things are hard for me to do. 1 2 3 4
40. I feel I need to punish myself for things I
have done or thought. 1 2 3 4
41. I am easy to love. 1 2 3 4
42. I often use alcohol and/or drugs. 1 2 3 4
43. I have thoughts of how to do myself in. 1 2 3 4
44. I find myself feeling depressed. 1 2 3 4
45. I think I have too much responsibility. 1 2 3 4
46. I sometimes get drunk or high alone. 1 2 3 4

APPENDIX B

Where To Get Help Information & Assistant Resources

American Association of Suicidology
2459 Ash
Denver, Colorado 80222
303/692-0985

Institute for Studies of Destructive Behaviors
1041 South Menlo Avenue
Los Angeles, California 90006
213/386-5111

National Center for Health Statistics
3700 East-West Highway, Room 157
Hyattsville, Maryland 20782
301/436-8500

National Institute of Mental Health
Public Inquiries Section, Room 15C05
5600 Fishers Lane
Rockville, Maryland 20857
301/443-4515

Youth Suicide National Center
West Coast Office
204 E. Second Avenue, Suite 203
San Mateo, California 94401
415/347-3961

S.P.A.R.E.
(Suicide Prevention Allied Regional Effort)
5100 South Picadilly
Aurora, Colorado 80015
303/680-5211

APPENDIX C

IMPORTANT FACTS

- Many adolescents who commit suicide hinted at their intent, so suicidal threats and attempts should always be treated seriously.
- Suicide is not inherited. However, research indicates that emotional disorders, such as depression, may be inherited. Therefore, children in some families may carry a higher likelihood of self-destruction than those in others.
- Suicidal people often are ambivalent about dying and frequently will call for help immediately following an attempted suicide.
- Many suicidal adolescents are depressed and distraught, most of them cannot be diagnosed as seriously mentally ill.
- Most adolescents who are suicidal are that way for only a brief period in their lives. If a young person who attempts suicide receives the proper assistance and support, he may never be suicidal again.
- Asking someone directly about a suspected suicidal intent often will lower their anxiety level by encouraging them to vent pent-up emotions, thus deterring the act of self-destruction.
- Suicide crosses all socioeconomic groups and no one class is more susceptible to it than another.
- The greatest danger of suicide exists during the first three months after an adolescent recovers from a deep depression.
- Most suicidal adolescents plan their self-destruction in advance.

*From Teens In Crisis (1989) by Thomas C. Barrett.

APPENDIX D

BASIC INTERVENTION STRATEGIES

- Don't be judgemental.
- Avoid saying, "No, you can't." No one can really prevent you from harming yourself if that is what you really want to do. But then again, no one can stop you a week from now or a month from now either. So, I suggest you postpone it until other solutions can be explored.
- Don't be sworn to secrecy.
- Get help from a mental health professional or agency.

*From Teens In Crisis (1989) by Thomas C. Barrett.

APPENDIX E

SIGNALS OF POTENTIAL BEHAVIOR

- Problems sleeping
- Loss of appetite
- Isolation and withdrawal
- Accident proneness
- Running away
- Dramatic changes in appearance
- Drug or alcohol abuse
- Agitation and aggression
- Talk about death
- Guilt feelings
- Hopelessness
- Giving away personal possessions
- Depression
- Short attention span
- Loss of interest in pleasurable activities
- Crying spells
- Self-criticism
- Sexual promiscuity
- A drop in school performance
- School attendance problems or dropping out
- Cult or gang membership
- Writings with messages of self-destruction
- Art work depicting violence, especially against self
- Euphoria following depression

STRESS-PRODUCING SITUATIONS

- Loss of someone significant through death, divorce, separation
- Loss of an important peer relationship
- Alienation
- Recent failure
- Family stress
- Moving

*From Teens In Crisis (1989) by Thomas C. Barrett

APPENDIX F

TEACHING COPING SKILLS IS THE KEY TO PREVENTING SUICIDE

PARENTS NEED TO BE AWARE OF:

Many parents, devastated by the suicide of a son or a daughter, recall certain behaviors that may have indicated a potential for the suicide. Others feel that there were no warning signs. No parent can fully know what to expect, but there are things a parent can do and know that might prove helpful in saving a child. Specifically:

- Extreme behavior patterns are not necessarily normal. Such behavior may be a sign that a child is disturbed.
- For teens who have limited coping skills, mild depression can turn to deeper depression with thoughts of suicide or another form of self-destruction.
- Be more aware of involvement with school, peers, and the community.
- Be empathetic when problems arise.
- Major changes in the family structure can be very difficult for an adolescent.
- If changes in a child's personality occur, seek qualified mental health advice for help.

*From Teen In Crisis (1989) by Thomas C. Barrett.

END

U.S. Dept. of Education

Office of Education
Research and
Improvement (OERI)

ERIC

Date Filmed

March 29, 1991